



**Europäisches
Patentamt**

**European
Patent Office**

**Office européen
des brevets**

PCT/EP200 4 / 0 1 2 9 5.7

16.11.04

Bescheinigung

Certificate

Attestation

Die angehefteten Unterla-
gen stimmen mit der
ursprünglich eingereichten
Fassung der auf dem näch-
sten Blatt bezeichneten
europäischen Patentanmel-
dung überein.

The attached documents
are exact copies of the
European patent application
described on the following
page, as originally filed.

Les documents fixés à
cette attestation sont
conformes à la version
initialement déposée de
la demande de brevet
européen spécifiée à la
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

03026779.3

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk

BEST AVAILABLE COPY



Europäisches
Patentamt

European
Patent Office

Office européen
des brevets

PCT/EP2004/012957

16.11.04

Anmeldung Nr:
Application no.: 03026779.3
Demande no:

Anmeldetag:
Date of filing: 21.11.03
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

Newron Pharmaceuticals S.p.A.
Via Ariosto, 21
20091 Bresso (MI)
ITALIE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se référer à la description.)

3-aminopoyrrolidone derivatives

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)
revendiquée(s)
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:


C07D207/00

Am Anmeldetag benannte Vertragsstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL
PT RO SE SI SK TR LI

3-AMINOPYRROLIDONE DERIVATIVES

This invention is related to 3-aminopyrrolidone derivatives and analogues of the following general formula I




I

and pharmaceutically acceptable salts thereof, that are active as sodium and/or calcium channel modulators and therefore useful in preventing, alleviating and curing a wide range of pathologies, including, but not limited to, neurological, psychiatric, cardiovascular, inflammatory, ophthalmic, urologic, metabolic and gastrointestinal diseases, where the above mechanisms have been described as playing a pathological role.

BACKGROUND OF THE INVENTION

Chemical background

WO97/05111 discloses N-(4-substituted-benzyl)-2-aminolactam derivatives of general formula II



II

wherein:

m is zero, 1, 2 or 3; n is zero, 1, 2 or 3; X is O, S, CH₂ or NH; each of R and R₁ independently is hydrogen, C₁-C₆ alkyl, halogen, hydroxy, C₁-C₄

alkoxy or trifluoromethyl; each of R_2 , R_3 and R_4 independently is hydrogen, C_1 - C_6 alkyl optionally substituted by a hydroxy group, or C_3 - C_7 cycloalkyl.

The compounds of the invention are active on the central nervous system (CNS) and can be used in therapy, for example as antiepileptics, in the treatment of Parkinson's disease and as neuroprotective agents, e.g. preventing or treating neuronal loss associated with stroke, ischemia, CNS trauma, hypoglycaemia or surgery and in treating or preventing neurodegenerative diseases such as Alzheimer's disease, amyotrophic lateral sclerosis, Down's syndrome or Huntington's disease; they can also be used as antidepressants, hypnotics and antispastic agents.

In WO 96/40679, WO 95/33719 and US 3,714,175 N-substituted pyrrolidone derivatives are claimed to be active as anticoagulants (Factor Xa), herbicides and muscle relaxants, respectively.

In EP 0362941, N-OH substituted pyrrolidone derivatives are claimed to be active as anticonvulsants, muscle relaxants and in anti-neurodegenerative disorders.

Biological background

It is well known that sodium channels play an important role in the neuronal network by transmitting electrical impulses rapidly throughout cells and cell networks, thereby coordinating higher processes ranging from locomotion to cognition. These channels are large transmembrane proteins, which are able to switch between different states to enable selective permeability for sodium ions. For this process an action potential is needed to depolarize the membrane, and hence these channels are voltage-gated. In the past few years a much better understanding of sodium channels and drugs interacting with them has been developed.

It has become clear that a number of drugs having an unknown mechanism of action actually act by modulating sodium channel conductance,

including local anesthetics, class I antiarrhythmics and anticonvulsants. Neuronal sodium channel blockers have found application with their use in the treatment of epilepsy (phenytoin and carbamazepine), bipolar disorder (lamotrigine), preventing neurodegeneration, and in reducing neuropathic pain. Various anti-epileptic drugs that stabilize neuronal excitability are effective in neuropathic pain (gabapentin).

In addition, an increase in sodium channel expression or activity has also been observed in several models of inflammatory pain, suggesting a role of sodium channels in inflammatory pain.

Calcium channels are membrane-spanning, multi-subunit proteins that allow controlled entry of calcium ions into cells from the extracellular fluid. Commonly, calcium channels are voltage dependent and are referred to as voltage sensitive calcium channels (VSCC). VSCCs are found throughout the mammalian nervous system, where they regulate such varied activities as cellular excitability, transmitter release, intracellular metabolism, neurosecretory activity and gene expression. All "excitable" cells in animals, such as neurons of the central nervous system (CNS), peripheral nerve cells, and muscle cells, including those of skeletal muscles, cardiac muscles and venous and arterial smooth muscles, have voltage dependent calcium channels. Calcium channels have a central role in regulating intracellular calcium ions levels that are important for cell viability and function. Intracellular calcium ion concentrations are implicated in a number of vital processes in animals, such as neurotransmitter release, muscle contraction, pacemaker activity, and secretion of hormones. It is believed that calcium channels are relevant in certain disease states. A number of compounds useful in treating various cardiovascular diseases in mammals, including humans, are thought to exert their beneficial effects by modulating functions of voltage dependant calcium channels present in cardiac and/or vascular smooth muscle.

Compounds with activity against calcium channels have also been implicated for the treatment of pain. In particular N-type calcium channels (Cav2.2), responsible for the regulation of neurotransmitters, are thought to play a significant role in nociceptive transmission, both due to their tissue
5 distribution as well as from the results of several pharmacological studies. This hypothesis has been validated in the clinic by Zinocotide, a peptide derived from the venom of the marine snail, *Conus Magus*. A limitation in the therapeutic use of this peptide is that it has to be administered intrathecally in humans (Bowersox S. S. and Luther R. *Toxicon*, 1998, 36, 11, 1651-1658).

10 All together these findings indicate that compounds with sodium and/or calcium channel blockade have a high therapeutic potential in preventing, alleviating and curing a wide range of pathologies, including neurological, psychiatric, cardiovascular, urologic, metabolic and gastrointestinal diseases, where the above mechanisms have been described as playing a pathological role.


15 There are many papers and patents which describe sodium channel and/or calcium channel modulators or antagonists for the treatment or modulation of a plethora of disorders, such as their use as local anesthetics, antiarrhythmics, antiemetics, antimanic depressants, agents for the treatment of unipolar depression, cardiovascular diseases, urinary incontinence, diarrhea,
20 inflammation, epilepsy, neurodegenerative conditions, nerve cell death, anticonvulsants, neuropathic pain, migraine, acute hyperalgesia and inflammation, renal disease, allergy, asthma, bronchospasm, dysmenorrhea, esophageal spasm, glaucoma, urinary tract disorders, gastrointestinal motility disorders, premature labour, obesity. A largely incomplete list is shown below.

25 An extensive and thorough prior art overview is reported in WO 03/057219 (and references therein); a further selection of prior art is reported in the following references: Alzheimer, C. *Adv. Exp. Med. Biol.* 2002, 513, 161-181; Vanegas, H.; Schaible, H. *Pain* 2000, 85, 9-18; U.S. Patent 5,051,403;

U.S. Patent 5,587,454; U.S. Patent 5,863,952; U.S. Patent 6,011,035;
 U.S. Patent 6,117,841; U.S. Patent 6,362,174; U.S. Patent 6,380,198;
 U.S. Patent 6,420,383; U.S. Patent 6,458,781; U.S. Patent 6,472,530;
 U.S. Patent 6,518,288; U.S. Patent 6,521,647; WO 97/10210; WO 03/018561.

5 DESCRIPTION OF THE INVENTION

The present invention relates to the use of compounds of formula I



I

10 wherein

m is an integer from 1 to 3

X is methylene, oxygen, sulphur or a NR⁶ group;

15 R¹ is a straight or branched C₁-C₈ alkyl or C₃-C₈ alkenylene or C₃-C₈ alkynylene chain, optionally substituted with CF₃, phenyl, phenoxy or naphthyl, the aromatic rings optionally substituted by one or more C₁-C₄ alkyl, halogens, trifluoromethyl, hydroxy or C₁-C₄ alkoxy groups;

R², R³ are independently hydrogen, a C₁-C₃ alkyl chain, halogen, trifluoromethyl, hydroxy or C₁-C₄ alkoxy groups;

20 R⁴, R⁵, R⁶ are independently hydrogen or C₁-C₆ alkyl;

and the pharmaceutically acceptable salts thereof for the preparation of a medicament having sodium and/or calcium channel modulating activity.

This invention is also related to the compounds of the general formula I

wherein m, X, R¹, R², R³, R⁴, R⁵ and R⁶ are as defined above with the

25 proviso that:

when R^1 is phenyl, benzyl, 2-phenethyl or 3-phenpropyl optionally and independently substituted on the phenyl ring by one or two C_1 - C_6 alkyl, halogen, hydroxy, C_1 - C_4 alkoxy or trifluoromethyl; X is oxygen, sulphur, methylene or -NH-, at least one of R^2 and R^3 is other than hydrogen. The
 5 disclaimed compounds are known from WO97/05111.

The optional substituents R^1 -X, R^2 and R^3 in the phenyl ring may be in any position.

The pharmaceutically acceptable salts of the compounds of formula I include acid addition salts with inorganic, e.g. hydrochloric, hydrobromic,
 10 sulphuric, and phosphoric acids or organic, e.g. acetic, propionic, benzoic, cinnamic, mandelic, salicylic, glycolic, lactic, oxalic, malic, maleic, malonic, fumaric, tartaric, citric, and the like.

Preferred use compounds of the invention are the compounds of formula I wherein m is 1 or 2, X is oxygen or methylene or NH or NCH_3 , R^1 is
 15 C_1 - C_8 alkyl chain, optionally substituted with CF_3 , phenyl or phenoxy group, where the aromatic ring in R^1 is optionally substituted by one or two halogen or methoxy or trifluoromethyl groups, R^2 and R^3 are hydrogen, methyl, methoxy, fluorine, chlorine or bromine, R^4 and R^5 are hydrogen or methyl, halogen is chlorine or fluorine.

20 Examples of specific use compounds of the invention are:

3-(4-Butyloxy-benzylamino)-pyrrolidin-2-one;

3-[4-(4-trifluorobutyloxy)-benzylamino]-pyrrolidin-2-one;

3-(4-Pentyloxy-benzylamino)-pyrrolidin-2-one;

3-[4-(5-trifluoropentyloxy)-benzylamino]-pyrrolidin-2-one;

25 3-(4-Phenylethyl-benzylamino)-pyrrolidin-2-one;

3-(4-Benzyloxy-benzylamino)-pyrrolidin-2-one;

3-(4-Phenylbutoxy-benzylamino)-pyrrolidin-2-one;

3-(4-Phenylpentyloxy-benzylamino)-pyrrolidin-2-one;

- 3-(4-Phenylallyloxy)-benzylamino-pyrrolidin-2-one;
3-(4-Phenoxyethoxy-benzylamino)-pyrrolidin-2-one;
3-[4-(Naphthalen-1-ylmethoxy)-benzylamino]-pyrrolidin-2-one;
3-(4-Pentyloxy-3-fluoro-benzylamino)-pyrrolidin-2-one;
5 3-(4-Pentyloxy-3-chloro-benzylamino)-pyrrolidin-2-one;
3-(4-Pentyloxy-3-bromo-benzylamino)-pyrrolidin-2-one;
3-(4-Pentyloxy-3-methoxy-benzylamino)-pyrrolidin-2-one;
3-(4-Pentyloxy-3-methyl-benzylamino)-pyrrolidin-2-one;
3-(4-Benzyloxy-3-fluoro-benzylamino)-pyrrolidin-2-one;
10 3-(4-Benzyloxy-3-bromo-benzylamino)-pyrrolidin-2-one;
3-(4-Benzyloxy-3-methoxy-benzylamino)-pyrrolidin-2-one;
3-(4-Benzyloxy-3-methyl-benzylamino)-pyrrolidin-2-one;
3-(4-Phenylpentoxy-2-chloro-benzylamino)-pyrrolidin-2-one;
3-(4-Phenylpentoxy-3-bromo-benzylamino)-pyrrolidin-2-one;
15 3-(4-Phenylpentoxy-3-methoxy-benzylamino)-pyrrolidin-2-one;
3-(4-Phenylpentoxy-3-methyl-benzylamino)-pyrrolidin-2-one;
3-(4-Phenylallyloxy-2-chloro-benzylamino)-pyrrolidin-2-one;
3-(4-Phenylallyloxy-3-fluoro-benzylamino)-pyrrolidin-2-one;
3-(4-Phenylallyloxy-3-bromo-benzylamino)-pyrrolidin-2-one;
20 3-(4-Phenylallyloxy-3-methoxy-benzylamino)-pyrrolidin-2-one;
3-(4-Phenylallyloxy-3-methyl-benzylamino)-pyrrolidin-2-one;
3-(4-Phenoxyethoxy-2-chloro-benzylamino)-pyrrolidin-2-one;
3-(4-Phenoxyethoxy-3-fluoro-benzylamino)-pyrrolidin-2-one;
3-(4-Phenoxyethoxy-3-bromo-benzylamino)-pyrrolidin-2-one;
25 3-(4-Phenoxyethoxy-3-methoxy-benzylamino)-pyrrolidin-2-one;
3-(4-Phenoxyethoxy-3-methyl-benzylamino)-pyrrolidin-2-one;
3-[4-(Naphthalen-1-ylmethoxy)-3-bromo-benzylamino]-pyrrolidin-2-one;
3-[4-(Naphthalen-1-ylmethoxy)-3-methoxy-benzylamino]-pyrrolidin-2-one;

- 3-[4-(Naphthalen-1-ylmethoxy)-3-methyl-benzylamino]-pyrrolidin-2-one;
3-(4-Pentyloxy-3-bromo-5-methoxy-benzylamino)-pyrrolidin-2-one;
3-(4-Pentyloxy-3,5-dimethoxy-benzylamino)-pyrrolidin-2-one;
3-(4-Pentyloxy-3,5-dimethyl-benzylamino)-pyrrolidin-2-one;
5 3-(4-Benzyloxy-3-bromo-5-methoxy-benzylamino)-pyrrolidin-2-one;
3-(4-Benzyloxy-3,5-dimethoxy-benzylamino)-pyrrolidin-2-one;
3-(4-Benzyloxy-3,5-dimethyl-benzylamino)-pyrrolidin-2-one;
3-(4-Phenylallyloxy-3-bromo-5-methoxy-benzylamino)-pyrrolidin-2-one;
3-(4-Phenylallyloxy-3,5-dimethoxy-benzylamino)-pyrrolidin-2-one;
10 3-(4-Phenylallyloxy-3,5-dimethyl-benzylamino)-pyrrolidin-2-one;
3-(4-Phenylpentoxy-3-bromo-5-methoxy-benzylamino)-pyrrolidin-2-one;
3-(4-Phenylpentoxy-3,5-dimethoxy-benzylamino)-pyrrolidin-2-one;
3-(4-Phenylpentoxy-3,5-dimethyl-benzylamino)-pyrrolidin-2-one;
3-(4-Phenoxyethoxy-3-bromo-5-methoxy-benzylamino)-pyrrolidin-2-one;
15 3-(4-Phenoxyethoxy-3,5-dimethoxy-benzylamino)-pyrrolidin-2-one;
3-(4-Phenoxyethoxy-3,5-dimethyl-benzylamino)-pyrrolidin-2-one;
3-[4-(Naphthalen-1-ylmethoxy)-2-chloro-5-methoxy-benzylamino]-
pyrrolidin-2-one;
3-[4-(Naphthalen-1-ylmethoxy)-3-fluoro-5-methoxy-benzylamino]-
20 pyrrolidin-2-one;
3-[4-(Naphthalen-1-ylmethoxy)-3-bromo-5-methoxy-benzylamino]-
pyrrolidin-2-one;
3-[4-(Naphthalen-1-ylmethoxy)-3,5-dimethoxy-benzylamino]-
pyrrolidin-2-one;
25 3-[4-(Naphthalen-1-ylmethoxy)-3,5-dimethyl-benzylamino]-pyrrolidin-
2-one;
3-[4-(2-Fluorobenzyloxy)-benzylamino]-pyrrolidin-2-one;
3-[4-(2-Fluorobenzyloxy)-benzylamino]-N-methylpyrrolidin-2-one;

- 3-[4-(2-triFluoromethyl-benzyloxy)-benzylamino]-pyrrolidin-2-one;
 3-[4-(2-Chlorobenzyloxy)-benzylamino]-pyrrolidin-2-one;
 3-[4-(2-Methoxybenzyloxy)-benzylamino]-pyrrolidin-2-one;
 3-[4-(3-Fluorobenzyloxy)-benzylamino]-pyrrolidin-2-one;
 5 3-[4-(3-Fluorobenzyloxy)-benzylamino]-N-methylpyrrolidin-2-one;
 3-{N-[4-(3-Fluorobenzyloxy)-benzyl]-N-methyl}-amino-pyrrolidin-2-one;
 3-[4-(3-triFluoromethyl-benzyloxy)-benzylamino]-pyrrolidin-2-one;
 3-[4-(3-Chlorobenzyloxy)-benzylamino]-pyrrolidin-2-one;
 3-[4-(3-Methoxybenzyloxy)-benzylamino]-pyrrolidin-2-one;
 10 3-[4-(3-Methoxybenzyloxy)-benzylamino]-N-methylpyrrolidin-2-one;
 3-[4-(4-Fluorobenzyloxy)-benzylamino]-pyrrolidin-2-one;
 3-[4-(4-Chlorobenzyloxy)-benzylamino]-pyrrolidin-2-one;
 3-[4-(4-Methoxybenzyloxy)-benzylamino]-pyrrolidin-2-one;
 3-[4-(4-triFluoromethyl-benzyloxy)-benzylamino]-pyrrolidin-2-one;
 15 3-[4-(2,3-diChlorobenzyloxy)-benzylamino]-pyrrolidin-2-one;
 3-[4-(3,4-diChlorobenzyloxy)-benzylamino]-pyrrolidin-2-one;
 3-[4-(3,4-diMethoxybenzyloxy)-benzylamino]-pyrrolidin-2-one;
 3-[4-(3,5-diMethoxybenzyloxy)-benzylamino]-pyrrolidin-2-one;
 3-[4-(3,5-diMethoxybenzyloxy)-benzylamino]-N-methylpyrrolidin-2-one;
 20 3-[4-(3,5-diMethoxyphenyl)-pentoxy]-benzylamino-pyrrolidin-2-one;
 3-[4-(2-Fluorobenzyloxy)-3-methyl-benzylamino]-pyrrolidin-2-one;
 3-[4-(2-triFluoromethyl-benzyloxy)-3-methyl-benzylamino]-pyrrolidin-
 2-one;
 3-[4-(3-Fluorobenzyloxy)-3-methyl-benzylamino]-pyrrolidin-2-one;
 25 3-{[4-(3-Fluorobenzyloxy)-3-methyl-benzyl]-N-methylamino}-
 pyrrolidin-2-one;
 3-[4-(3-triFluoromethyl-benzyloxy)-3-methyl-benzylamino]-pyrrolidin-
 2-one;

- 3-[4-(3-Chlorobenzoyloxy)-3-methyl-benzylamino]-pyrrolidin-2-one;
3-{[4-(3-Chlorobenzoyloxy)-3-methyl-benzyl]-N-methylamino}-
pyrrolidin-2-one;
- 5 3-[4-(3-Bromobenzoyloxy)-3-methyl-benzylamino]-pyrrolidin-2-one;
3-{[4-(3-Bromobenzoyloxy)-3-methyl-benzyl]-N-methylamino}-
pyrrolidin-2-one;
- 3-[4-(4-triFluoromethyl-benzoyloxy)-2-chloro-benzylamino]-pyrrolidin-
2-one;
- 10 3-[4-(4-Fluorobenzoyloxy)-3-methyl-benzylamino]-pyrrolidin-2-one;
3-[4-(4-triFluoromethyl-benzoyloxy)-3-fluoro-benzylamino]-pyrrolidin-
2-one;
- 3-[4-(4-triFluoromethyl-benzoyloxy)-3-bromo-benzylamino]-pyrrolidin-
2-one;
- 3-[4-(4-triFluoromethyl-benzoyloxy)-3-methoxy-benzylamino]-
15 pyrrolidin-2-one;
- 3-[4-(4-triFluoromethyl-benzoyloxy)-3-methyl-benzylamino]-pyrrolidin-
2-one;
- 3-[4-(4-Chlorobenzoyloxy)-3-methyl-benzylamino]-pyrrolidin-2-one;
- 3-[4-(4-triFluoromethyl-benzoyloxy)-3-bromo-5-methoxy-benzylamino]-
20 pyrrolidin-2-one;
- 3-[4-(4-triFluoromethyl-benzoyloxy)-3,5-dimethoxy-benzylamino]-
pyrrolidin-2-one;
- 3-[4-(4-triFluoromethyl-benzoyloxy)-3,5-dimethyl-benzylamino]-
pyrrolidin-2-one;
- 25 3-[4-(3,4-diChlorobenzoyloxy)-2-chloro-benzylamino]-pyrrolidin-2-one;
3-[4-(3,4-diChlorobenzoyloxy)-3-fluoro-benzylamino]-pyrrolidin-2-one;
3-[4-(3,4-diChlorobenzoyloxy)-3-bromo-benzylamino]-pyrrolidin-2-one;
3-[4-(3,4-diChlorobenzoyloxy)-3-methoxy-benzylamino]-pyrrolidin-2-one;


- 3-[4-(3,4-diChlorobenzyloxy)-3-methyl-benzylamino]-pyrrolidin-2-one;
3-[4-(3,5-diMethoxybenzyloxy)-2-chloro-benzylamino]-pyrrolidin-2-one;
3-[4-(3,5-diMethoxybenzyloxy)-3-fluoro-benzylamino]-pyrrolidin-2-one;
3-[4-(3,5-diMethoxybenzyloxy)-3-bromo-benzylamino]-pyrrolidin-2-one;
5 3-[4-(3,5-diMethoxybenzyloxy)-3-methoxy-benzylamino]-pyrrolidin-2-one;
3-[4-(3,5-diMethoxybenzyloxy)-3-methyl-benzylamino]-pyrrolidin-2-one;
3-[4-(3,4-diChlorobenzyloxy)-3,5-dimethoxy-benzylamino]-pyrrolidin-
2-one;
3-[4-(3,4-diChlorobenzyloxy)-3,5-dimethyl-benzylamino]-pyrrolidin-2-one;
10 3-[4-(3,5-diChlorobenzyloxy)-3-bromo-5-methoxy-benzylamino]-
pyrrolidin-2-one;
3-[4-(3,5-diMethoxybenzyloxy)-3-bromo-5-methoxy-benzylamino]-
pyrrolidin-2-one;
3-[4-(3,5-diMethoxybenzyloxy)-3,5-dimethoxy-benzylamino]-
15 pyrrolidin-2-one;
3-[4-(3,5-diMethoxyphenyl)-allyloxy)-3,5-dimethoxy-benzylamino]-
pyrrolidin-2-one.
3-(4-Benzyloxy-benzylamino)-piperidin-2-one;
3-(4-Benzyloxy-benzylamino)-azepan-2-one;
20 3-[4-(2-Fluorobenzyloxy)-benzylamino]-piperidin-2-one;
3-[4-(2-Fluorobenzyloxy)-benzylamino]-azepan-2-one;
3-[4-(2-Chlorobenzyloxy)-benzylamino]-piperidin-2-one;
3-[4-(2-Chlorobenzyloxy)-benzylamino]-azepan-2-one;
3-[4-(3-Fluorobenzyloxy)-benzylamino]-piperidin-2-one;
25 3-[4-(3-Fluorobenzyloxy)-benzylamino]-azepan-2-one;
3-[4-(4-Fluorobenzyloxy)-benzylamino]-piperidin-2-one;
3-[4-(4-Fluorobenzyloxy)-benzylamino]-azepan-2-one;
3-[4-(2-Chlorobenzylamino)-benzylamino]-piperidin-2-one;

12

3-[4-(2-Chlorobenzylamino)-benzylamino]-azepan-2-one;
 3-{4-[(2-Chlorobenzyl)methylamino]-benzylamino}-piperidin-2-one;
 3-{4-[(2-Chlorobenzyl)methylamino]-benzylamino}-azepan-2-one;
 or pharmaceutically acceptable salts thereof.

5 The compounds of the invention and the salts thereof can be obtained
 by a process comprising:


a) reaction of compounds of formula II



II

10

wherein R¹, R², R³ and X are as defined above
 with compounds of formula III, in the presence of a reducing agent




III

15

wherein m and R⁵ are as defined previously thus obtaining a compound
 of formula I; or

b) reaction of compounds of formula IV



IV


20

13

wherein R^1 , R^2 , R^3 and X are as defined above and Y is a halogen atom or a O-EWG group, where the EWG means an electron withdrawing group, like e.g. mesyl, tosyl or trifluoroacetyl groups, able to transform the oxygen which they are linked to, in a good leaving group

5 with compounds of formula III thus obtaining a compound of formula I; or

c) reacting of a compound of formula V



V

10 wherein R^1 , R^2 , R^3 , R^5 , X and m are as defined above, with compounds of formula VI or VII



VI



VII

15 wherein Y is as defined above; R^4 is as above defined and R^7 is hydrogen or C_1-C_5 alkyl; and, if desired, converting a compound of the invention into another compound of the invention and/or, if desired, converting a compound of the invention into a pharmaceutically acceptable salt and/or, if desired, converting a salt into a free compound and/or, if desired, separating a mixture of isomers of compounds of the invention into a single isomer.

20 Compounds II, III, IV, VI and VII are commercially available compounds or are prepared from commercially available compounds using well-known methods.

The reactions of compounds of formula II with compounds of formula III and of compounds of formula V with compounds of formula VII to give the compounds of formula I is a reductive amination reaction which can be carried out according to known methods. According to a preferred embodiment of the invention, it may be performed under nitrogen atmosphere, in a suitable organic solvent, such as an alcohol, e.g. a lower alkanol, in particular methanol, or in acetonitrile, or in tetrahydrofuran, at a temperature ranging from about 0°C to about 70°C, in the presence of a reducing agent, the most appropriate being sodium borohydride or sodium cyanoborohydride. Occasionally Titanium IV isopropylate and molecular sieves can be added to the reaction mixture for facilitating the reaction.

In a compound of formula IV and VI the halogen is preferably bromine or iodine. The alkylation reactions of a compound of formula IV with a compound of formula III and of a compound of formula V with a compound of formula VI can be carried out in a suitable organic solvent, such as an alcohol, e.g. methanol, ethanol or isopropanol, in particular in ethanol, at a temperature ranging from about 0°C to about 50°C.

When in the compounds of the present invention and in the intermediate-products thereof, groups are present, which need to be protected before submitting them to the above illustrated reactions, they may be protected before being reacted and then deprotected according to well-known methods.

PHARMACOLOGY

The compounds of the invention display affinities for the calcium and/or sodium channel binding sites as demonstrated using selective radioligands in *in vitro* binding studies.

The compounds according to the invention are blockers of the voltage dependent sodium channels and/or calcium channels. These compounds

therefore displace 3H-batrachotoxin (BTX) with a high affinity from the binding site on the sodium channel, with IC_{50} in the low μM or sub μM range. Similarly the compounds displace 3H-nitrendipine from the binding site in the calcium channel, with IC_{50} in the low μM or most usually in the sub μM range as well as inhibiting calcium influx induced through calcium channels via cellular depolarisation.

Such substances exhibit "use-dependency" when the sodium channels are blocked i.e. maximum blockage of the sodium channels is only achieved after repeated stimulation of the sodium channel. Consequently, the substances preferably bind to sodium channels which are multiply activated. As a result the substances are capable of activity preferentially in those regions of the body which are pathologically over-stimulated, as illustrated by patch-clamp experiments (W.A. Catterall, Trends Pharmacol. Sci., 8, 57-65; 1987) which show that the compounds according to the invention block the electrically stimulated sodium channel in a "use-dependent" manner.

As a consequence of these mechanisms the compounds of the invention are active *in vivo* when orally administered in the range of 0.1 to 100 mg/kg in animal models such as the formalin model of persistent pain and the carragenan model of inflammation.

In view of the above described mechanisms of action, the compounds of the present invention are useful in the treatment or prevention of neuropathic pain. Neuropathic pain syndromes include, and are not limited to: diabetic neuropathy; sciatica; non-specific lower back pain; multiple sclerosis pain; fibromyalgia; HIV-related neuropathy; neuralgia, such as post-herpetic neuralgia and trigeminal neuralgia; and pain resulting from physical trauma, amputation, cancer, toxins or chronic inflammatory conditions.

Compounds of the invention are also useful for the treatment of chronic pain. Chronic pain includes, and is not limited to, chronic pain caused by

inflammation or an inflammatory-related condition, osteoarthritis, rheumatoid arthritis or as sequela to disease, acute injury or trauma and includes upper back pain or lower back pain (resulting from systematic, regional or primary spine disease (such as radiculopathy), bone pain (due to
5 osteoarthritis, osteoporosis, bone metastasis or unknown reasons), pelvic pain, spinal cord injury-associated pain, cardiac chest pain, non-cardiac chest pain, central post-stroke pain, myofascial pain, cancer pain, AIDS pain, sickle cell pain, geriatric pain or pain caused by headache, temporomandibular joint syndrome, gout, fibrosis or thoracic outlet syndromes.

10 Compounds of the invention are also useful in the treatment of acute pain (caused by acute injury, illness, sports-medicine injuries, carpal tunnel syndrome, burns, musculoskeletal sprains and strains, musculotendinous strain, cervicobrachial pain syndromes, dyspepsis, gastric ulcer, duodenal ulcer, dysmenorrhea, endometriosis or surgery (such as open heart or bypass
15 surgery), post operative pain, kidney stone pain, gallbladder pain, gallstone pain, obstetric pain or dental pain.

Compounds of the invention are also useful in the treatment of migraine, such as tension type headache, transformed migraine or evolutive
headache, cluster headache, as well as secondary headache disorders, such as
20 the ones derived from infections, metabolic disorders or other systemic illnesses and other acute headaches, paroxysmal hemicrania and the like, resulting from a worsening of the above mentioned primary and secondary headaches.

Compounds of the invention are also useful for the treatment of
25 neurological conditions and cognitive disorders. Neurological conditions include, and are not limited to, conditions such as degenerative dementia (including senile dementia, and Pick's diseases) dementia associated with intracranial space occupying lesions, infections and related conditions

(including HIV infection), metabolism, toxins, anoxia and vitamin deficiency; and mild cognitive impairment associated with aging, particularly age associated memory impairment, movement disorders (postencephalitic parkinsonism, progressive supranuclear palsy, corticobasal degeneration),
5 narcolepsy, deficit and hyperactivity disorders (ADHD).

Compounds of the invention are also useful for the treatment of psychiatric disorders. Psychiatric disorders include, and are not limited to, bipolar disorder (such as bipolar disorder type I, bipolar disorder type II), cyclothymic disorder, rapid cycling, ultradian cycling, mania, hypomania,
10 schizophrenia, schizophreniform disorders, schizoaffective disorders, delusional disorders, brief psychotic disorders, shared psychotic disorders, psychotic disorder due to a general medical condition, substance-induced psychotic disorders or a psychotic disorder not otherwise specified, anxiety disorders and moreover in smoke and drug addiction.

15 Compounds of the invention are also useful in the treatment of peripheral diseases such as tinnitus, muscular sclerosis, and other disorders, including, and not limited to cardiovascular diseases (such as cardiac arrhythmia, cardiac infarction or angina pectoris, hypertension, hypoxia, cardiac ischemia) endocrine disorders (such as acromegaly or diabetes insipidus), diseases in
20 which the pathophysiology of the disorder involves excessive or hypersecretory or otherwise inappropriate cellular secretion of an endogeneous substance (such as catecholamine, a hormone or a growth factor).

Compounds of the invention are also useful in the treatment of liver disease, such as inflammatory liver disease, for example chronic viral hepatitis
25 B, chronic viral hepatitis C, alcoholic liver injury, primary biliary cirrhosis, autoimmune hepatitis, non-alcoholic steatohepatitis and liver transplant rejection.

Compounds of the invention inhibit inflammatory processes affecting all

body systems. Therefore are useful in the treatment of inflammatory processes of the muscular-skeletal system of which the following is a list of examples but it is not comprehensive of all target disorders: arthritic conditions such as alkylosing spondylitis, cervical arthritis, fibromyalgia, gut, juvenile rheumatoid arthritis, lumbosacral arthritis, osteoarthritis, osteoporosis, psoriatic arthritis, 5 rheumatic disease; disorders affecting skin and releted tissues: eczema, psoriasis, dermatitis and inflammatory conditions such as sunburn; disorders of the respiratory system: asthma, allergic rhinitis and respiratory distress syndrome, lung disorders in which inflammation is involved such as asthma and 10 bronchitis; chronic obstructive pulmonary disease; disorders of the immune and endocrinological systems: periarteritis nodosa, thyroiditis, aplastic anaemia, sclerodoma, myasthenia gravis, multiple sclerosis, sarcoidosis, nephritic syndrome, Bechet's syndrome, polymyositis, gingivitis.

Compounds of the invention are also useful in the treatment of 15 gastrointestinal (GI) tract disorders such as inflammatory bowel disorders including but not limited to ulcerative colitis, Crohn's disease, ileitis, proctitis, celiac disease, enteropathies, microscopic or collagenous colitis, eosinophilic gastroenteritis, or pouchitis resulting after proctocolectomy and post ileonatal anastomosis, and irritable bowel syndrome including any disorders associated 20 with abdominal pain and/or abdominal discomfort such as pylorospasm, nervous indigestion, spastic colon, spastic colitis, spastic bowel, intestinal neurosis, functional colitis, mucous colitis, laxative colitis and functional dyspepsia; but also for treatment of atrophic gastritis, gastritis varialoforme, ulcerative colitis, peptic ulceration, pyresis, and other damage to the GI tract, for example, by 25 *Helicobacter pylori*, gastroesophageal reflux disease, gastroparesis, such as dlabetic gastroparesis; and other functional bowel disorders, such as non-ulcerative dyspepsia (NUD); emesis, diarrhea, and visceral inflammation.

Compounds of the invention are also useful in the treatment of disorders

of the genito-urinary tract such as overactive bladder, prostatitis (chronic bacterial and chronic non-bacterial prostatitis), prostatic dysuria, interstitial cystitis, urinary incontinence and benign prostatic hyperplasia, annexites, pelvic inflammation, bartolinitis and vaginitis.

- 5 Compounds of the invention are also useful in the treatment of ophthalmic diseases such as retinitis, retinopathies, uveitis and acute injury to the eye tissue, macular degeneration or glaucoma, conjunctivitis.

Compounds of the invention are also useful in the treatment of obesity.

- 10 Compounds of the invention are also useful for the treatment of all other conditions mediated by the inhibition of voltage gated sodium channels and/or voltage gated calcium channels.

- It will be appreciated that the compounds of the invention may advantageously be used in conjunction with one or more other therapeutic agents. Examples of suitable agents for adjunctive therapy include a 5HT_{1B/1D} agonist, such as a triptan (e.g. sumatriptan or naratriptan); an adenosine A₁ agonist; an EP ligand; an NMDA modulator, such as a glycine antagonist; a substance P antagonist (e.g. an NK1 antagonist); a cannabinoid; acetaminophen or phenacetin; a 5-lipoxygenase inhibitor; a leukotriene receptor antagonist; a DMARD (e.g. methotrexate); gabapentin and related compounds; a tricyclic antidepressant (e.g. amitriptyline); a neurone stabilising antiepileptic drug; a monoaminergic uptake inhibitor (e.g. venlafaxine); a matrix metalloproteinase inhibitor; a nitric oxide synthase (NOS) inhibitor, such as an iNOS or an nNOS inhibitor; an inhibitor of the release, or action, of tumor necrosis factor alpha; an antibody therapy, such as monoclonal antibody therapy; an antiviral agent, such as a nucleoside inhibitor (e.g. lamivudine) or an immune system modulator (e.g. interferon); an analgesic, such as a cyclooxygenase-2 inhibitor; a local anaesthetic; a stimulant, including caffeine; an H₂-antagonist (e.g. ranitidine); a proton pump inhibitor (e.g. omeprazole); an antacid (e.g.
- 15
20
25

aluminium or magnesium hydroxide; an antifatulent (e.g. semethicone); a decongestant (e.g. phenylephrine, phenylpropanolamine, pseudoephedrine, oxymetazoline, epinephrine, naphazoline, xylometazoline, propylhexedrine, or levo-desoxyephedrine, naphazoline, xylometazoline, propylhexedrine, or levo-desoxyephedrine); antitussive (e.g. codeine, hydrocodone, carmiphen, carbetapentane, or dexamethorphan); a diuretic; or a sedating or non-sedating antihistamine. It is to be understood that the present invention covers the use of a compound of formula (I) or a pharmaceutically acceptable salt thereof in combination with one or more therapeutic agents.

10 The compounds of the present invention are useful in human and veterinary medicine. It is to be understood that reference to treatment includes both treatments of established symptoms and prophylactic treatment, unless explicitly stated otherwise.

15 The 3-aminopyrrolidone derivatives of formula I as above defined can be administered as the "active ingredient" of a pharmaceutically acceptable composition which can be prepared by conventional procedures, for instance by mixing the active ingredient with pharmaceutically acceptable, therapeutically inert organic and/or inorganic carrier materials.

20 The composition comprising the above defined 3-aminopyrrolidone derivatives can be administered in a variety forms, e.g. orally, in the form of tablets, troches, capsules, sugar or film coated tablets, liquid solutions, emulsions or suspensions; rectally, in the form of suppositories; parenterally, e.g. by intramuscular or intravenous injection or infusion; and transdermally.

25 Suitable pharmaceutically acceptable, therapeutically inert organic and/or inorganic carrier materials useful in the preparation of such composition include, for example, water, gelatin, gum arabic, lactose, starch, cellulose, magnesium stearate, talc, vegetable oils, polyalkyleneglycols and the like. The composition comprising the 3-aminopyrrolidone derivatives of

formula I as above defined can be sterilized and may contain further well known components, such as, for example, preservatives, stabilizers, wetting or emulsifying agents, e.g. paraffin oil, mannide monooleate, salts to adjust osmotic pressure, buffers and the like.

5 For example, the solid oral forms may contain, together with the active ingredient, diluents, e.g. lactose, dextrose, saccharose, cellulose, corn starch or potato starch; lubricants, e.g. silica, talc, stearic acid, magnesium or calcium stearate, and/or polyethylene glycols; binding agents, e.g. starches, arabic gums, gelatin, methylcellulose, carboxymethylcellulose or polyvinyl
10 pyrrolidone; disgregating agents, e.g. a starch, alginic acid, alginates or sodium starch glycolate; effervescing mixtures; dyestuffs; sweeteners; wetting agents such as lecithin, polysorbates, laurylsulphates; and, in general, non-toxic and pharmacologically inactive substances used in pharmaceutical formulations. Said pharmaceutical preparations may be manufactured in
15 known manner, for example, by means of mixing, granulating, tableting, sugar-coating, or film-coating processes.

The oral formulations comprise sustained release formulations that can be prepared in conventional manner, for instance by applying an enteric coating to tablets and granules.

20 The liquid dispersion for oral administration may be e.g. syrups, emulsions and suspensions.

The syrups may contain as carrier, for example, saccharose or saccharose with glycerine and/or mannitol and/or sorbitol.

25 Suspensions and emulsions may contain as a carrier, for example, a natural gum, agar, sodium alginate, pectin, methylcellulose, carboxymethylcellulose, or polyvinyl alcohol. The suspensions or solutions for intramuscular injections may contain, together with the active compound, a pharmaceutically acceptable carrier, e.g. sterile water, olive oil, ethyl oleate, glycols, e.g.

propylene glycol, and, if desired, a suitable amount of lidocaine hydrochloride. The solutions for intravenous injections or infusion may contain as carrier, for example, sterile water or preferably they may be in the form of sterile, aqueous, isotonic saline solutions.

5 The suppositories may contain, together with the active ingredient, a pharmaceutically acceptable carrier, e.g. cocoa butter, polyethylene glycol, a polyoxyethylene sorbitan fatty acid ester surfactants or lecithin.

Suitable treatment is given 1, 2 or 3 times daily, depending upon clearance rate. Accordingly, the desired dose may be presented in a single
10 dose or as divided doses administered at appropriate intervals, for example two to four or more sub-doses per day.

The pharmaceutical compositions comprising the 3-aminopyrrolidone derivatives of formula I as above defined will contain, per dosage unit, e.g., capsule, tablet, powder injection, teaspoonful, suppository and the like from about
15 0.1 to about 500 mg of the active ingredient most preferably from 1 to 10 mg.

Optimal therapeutically effective doses to be administered may be readily determined by those skilled in the art and will vary, basically, with the strength of the preparation, with the mode of administration and with the advancement of the condition or disorder treated. In addition, factors
20 associated with the particular subject being treated, including subject age, weight, diet and time of administration, will result in the need to adjust the dose to an appropriate therapeutically effective level.

The following examples further illustrate the invention.

Example 1

25 (S)-3-[4-(3-Fluorobenzyloxy)-benzylamino]-pyrrolidin-2-one

To a 2M solution of (S)-3-aminopyrrolidin-2-one (1.47 g, 14.6 mmol), prepared as described in Example 5, in anhydrous methanol 2.0 g of 3 Å molecular sieves and 0.77 g NaBH₃CN (12 mmol) were added; after 10

23

minutes, 3.4 g (14.8 mmol) of 4-(3-fluorobenzyloxy)benzaldehyde in 40 ml of anhydrous methanol were added. The reaction was carried out for 3 hours, then the mixture filtered, the solution was evaporated yielding a residue which was directly flash-chromatographed on silica gel (eluant: $\text{CHCl}_3:\text{CH}_3\text{OH}:\text{NH}_3$ (30%) 97:3:0.3, v:v:v) to give a white solid (3.4 g; 74%). The free base thus obtained was treated with a stoichiometric amount of methanesulfonic acid to give the title compound m.p. 140.5-143.5°C.

MS (ESI Pos Spray 3.5 kV; Skimmer 20 V; Probe 250°C): 315 $[\text{MH}^+]$

Example 2

10 (S)-3-{N-[4-(3-Fluorobenzyloxy)-benzyl]-N-methyl}-amino-pyrrolidin-2-one

1.0 g (3.2 mmol) of (S)-3-[4-(3-fluorobenzyloxy)-benzylamino]-pyrrolidin-2-one was dissolved in 50 ml acetonitrile under inert conditions. 1.6 ml (20 mmol) of 37% formaldehyde and 0.29 g (4.6 mmol) of sodium cyanoborohydride were added at room temperature. After 20 minutes glacial acetic acid was added until reaching a neutral pH value. The mixture was stirred for 40 minutes and then evaporated to dryness. After adding 40 ml of 2M potassium hydroxide the product was extracted with ethyl acetate, the organic layer was washed first with a 0.5 M KOH solution, then with water and brine. The organic layer was dried over Na_2SO_4 , then filtered and evaporated to obtain a residue which was flash-chromatographed on silica gel (eluant: $\text{CHCl}_3:\text{CH}_3\text{OH}:\text{NH}_3$ (30%), 200:3:0.2, v:v:v) to give 0.73 g (70%) of a white solid m.p. 89-92°C.

MS (ESI Pos Spray 3.5 kV; Skimmer 20 V; Probe 250°C): 329 $[\text{MH}^+]$

25 Example 3

(S)-3-[3,5-diMethyl-4-(4-trifluoromethyl-benzyloxy)-benzylamino]-pyrrolidin-2-one

2M solution of (S)-3-aminopyrrolidin-2-one (650 mg, 6.5 mmol),

prepared as described in Example 5, in dry THF was added to a 1M solution of 3,5-dimethyl-4-(4-trifluoromethyl-benzyloxy)-benzaldehyde (2.2 g, 7.15 mmol), prepared as described in Example 6, in dry THF. To this mixture, a 2M solution of Ti(O-*i*Pr)₄ (2.77 g, 9.75 mmol) in dry THF was added dropwise and the reaction mixture was stirred 12 hours at room temperature under nitrogen.

A 0.75M solution of sodium borohydride (690 mg, 18.2 mmol) in absolute ethanol was added and the resulting mixture was heated at 70°C for 6 hours. After cooling, water (8 ml) was added and the resulting white precipitate was removed by filtration. The crude compound was purified using SCX cartridge (Strong Cation eXchange resin). The pure product was recovered by eluting with 3% ammonia solution in methanol. The title compound was obtained (2.24 g) after evaporating the solvent under vacuum with a yield of 88%.

MS (ESI Pos Spray 3.5 kV; Skimmer 20 V; Probe 250°C): 393 [MH⁺]
¹H-NMR (DMSO-*d*₆) δ: 7.7 (m, 5H), 7.0 (s, 2H), 4.9 (s, 2H), 3.68 (s, 2H), 3.15 (m, 3H), 2.22 (m, 7H), 1.7 (m, 1H).

Example 4

(S)-3-[4-(3,4-Dichloro-benzyloxy)-3-methyl-benzylamino]-pyrrolidin-2-one

2M solution of (S)-3-amino-pyrrolidin-2-one (650 mg, 6.5 mmol), prepared as described in Example 5, in dry THF was added to a 1M solution of 4-(3,4-dichloro-benzyloxy)-3-methyl-benzaldehyde (2.1 g, 7.15 mmol), prepared as described in Example 7, in dry THF. To this mixture, a 2M solution of Ti(O-*i*Pr)₄ (2.77 g, 9.75 mmol) in dry THF was added dropwise and the reaction mixture was stirred 12 hours at room temperature under nitrogen.

A 0.75M solution of sodium borohydride (690 mg, 18.2 mmol) in

absolute ethanol was added and the resulting mixture was heated at 70°C for 6 hours. After cooling, water (8 ml) was added and the resulting white precipitate was removed by filtration. The crude compound was purified using SCX cartridge (Strong Cation eXchange resin). The pure product was recovered by eluting with 3% ammonia solution in methanol. The title compound was obtained (2.18 g) after evaporating the solvent under vacuum with a yield 89%.

MS (ESI Pos Spray 3.5 kV; Skimmer 20 V; Probe 250°C): 378.9 [MH⁺]

¹H-NMR (DMSO-d₆) δ: 7.72-7.68 (m, 3H), 7.45 (dd, 1H), 7.1 (m, 2H), 6.9 (dd, 1H), 5.15 (s, 2H), 3.65 (s, 2H), 3.15 (m, 3H), 2.20 (m, 4H), 1.68 (m, 1H).

Example 5

(S)-3-ammino-pyrrolidin-2-one

5M solution of hexamethyldisilazane (108 ml, 523 mmol) in CH₃CN was added dropwise to a solution of (S)-2,4-diamino-butyric acid (10 g, 52.3 mmol) in CH₃CN (150 ml) at room temperature. The resulting mixture was refluxed for 30 hours. The crude reaction was poured into cold MeOH (400 ml), stirred at room temperature for 30 minutes and then evaporated under vacuum. The resulting solid was dissolved in CH₂Cl₂ (700 ml) and the insoluble residue was removed by filtration under vacuum.

The title compound was obtained (3.13 g) after evaporating the solution to dryness in a 60% yield.

Example 6

3,5-Dimethyl-4-(4-trifluoromethyl-benzyloxy)-benzaldehyde

0.5M solution of 1-bromomethyl-4-trifluoromethyl-benzene (1.9 g, 8.0 mmol) in DMF was added dropwise to a suspension of 4-hydroxy-3,5-dimethyl-benzaldehyde (930 mg, 7.3 mmol), K₂CO₃ (1.51 g, 11 mmol) and KI (120 mg, 0.73 mmol) in DMF (100 ml). The reaction mixture was stirred at

90°C overnight. After cooling, the solid residue was filtered off and the solvent was evaporated under vacuum. The residue was dissolved in ethyl acetate and the organic layer washed twice with NaOH 1M, dried over Na₂SO₄ and evaporated to dryness. The residue was purified on silica gel, obtaining
5 2.24 g of the title compound in a quantitative yield.

Example 7

4-(3,4-diChloro-benzyloxy)-3-methyl-benzaldehyde

0.5M solution of 4-chloromethyl-1,2-dichloro-benzene (1.55 g, 8.0 mmol) in DMF was added dropwise to a suspension of 4-hydroxy-3-methyl-
10 benzaldehyde (990 mg, 7.3 mmol), K₂CO₃ (1.51 g, 11 mmol) and KI (120 mg, 0.73 mmol) in DMF (100 ml). The reaction mixture was treated as described in Example 6. The residue was purified on silica gel, obtaining 2.15 g of the title compound in a quantitative yield.

Example 8

***In vitro* binding of ³H-batrachotoxin in rat brain membranes**

Membrane preparations (P2 fraction) Male Wistar rats (Harlan, Italy – 175-200 g) were sacrificed under light anaesthesia and brains were rapidly removed and cortex was dissected out, homogenized in 10 vol. of ice-cold 0.25 M sucrose buffer (50 mM Tris-HCl, pH 7.4). The crude homogenate was
20 centrifuged at 3250 rpm for 10 min and the supernatant recovered. The pellet was homogenized and centrifuged again and the two supernatants were pooled and centrifuged at 14750 rpm for 10 min, +4°C. The resulting pellet was stored at -20°C until use.

Binding assay. The pellet was resuspended in 50 mM Hepes buffer, pH
25 7.4 containing 0.8 mM MgSO₄, 5.4 mM KCl, 5.5 mM glucose and 130 mM choline using Polytron PT10. The binding assay was carried out in 0.25 ml final volume containing 50 µl of membrane preparation (ca 200 µg of protein), 50 µl of ³H-batrachotoxin ligand (10 nM), 50 µl of TTX (1 µM), 50 µl of

scorpion toxin (37.5 µg/ml) and 50 µl of test compound or buffer or 300 µM of veratridine to determine non specific binding. The binding assay was performed at 37°C for 30 min and stopped by rapid filtration under vacuum through Whatman GF/B glass fiber filters. Filters (pre-soaked with polyethylenimine 0.1%) were washed with 3x5 ml of ice-cold buffer and placed in picovials containing scintillation cocktail (Filter Count, Packard). Radioactivity bound was measured by liquid scintillation spectrometry at 45% efficiency.

Data analysis. The IC₅₀ was calculated from displacement curves by computer program LIGAND (McPherson, J. Pharmacol. Methods, 14, 213. 1985). The displacement curves were obtained using at least 9 concentrations, each in duplicate, covering a 100000 fold range.

Example 9

In vitro binding of ³H-nitrendipine in rat brain membranes

Membrane preparations Male Wistar rats (Harlan, Italy – 175-200 g) were sacrificed under light anaesthesia and brains were rapidly removed and cortex was dissected out, homogenized in 10 vol. of ice-cold 50 mM Tris·HCl, pH 7.7 using Polytron PT10. The crude homogenate was centrifuged at 50000xg for 10 min. The pellet was homogenized and centrifuged twice in fresh buffer at 50000xg for 10 min, +4°C. The resulting pellet was stored at -20°C until use.

Binding assay. The pellet was resuspended in 50 mM Tris·HCl, pH 7.7 using Polytron PT10. The binding assay was carried out in 1 ml final volume containing 900 µl of membrane preparation (ca 700 µg of protein), 50 µl of ³H-nitrendipine (0.15 nM), and 50 µl of test compound or buffer or 1 µM of nifedipine to determine non specific binding. The binding assay was performed at 25°C for 45 min and stopped by rapid filtration under vacuum through Whatman GF/B glass fiber filters. Filters were washed with 3x5 ml of ice-cold buffer and placed in picovials containing scintillation cocktail (Filter

Count, Packard). Radioactivity bound was measured by liquid scintillation spectrometry at 45% efficiency.

Data analysis. The IC_{50} was calculated from displacement curves by computer program LIGAND (McPherson, J. Pharmacol. Methods, 14, 213, 1985). The displacement curves were obtained using at least 9 concentrations, each in duplicate, covering a 100000 fold range.

Example 10

Calcium influx assay

IMR32 human neuroblastoma cells constitutively possess both L and N type channels. Under differentiating conditions, IMR32 preferentially express on the membrane surface N- type calcium channels. The remaining L-type calcium channels were blocked using the selective L type blocker, nifedipine. In these experimental conditions, only N type channels can be detected.

IMR32 cells were differentiated using 1mM dibutyryl-cAMP and 2.5 μ M bromodeoxyuridine for 8 days (4 times) in 225 cm² flask, then detached, seeded at 200,000 cells/well on 96 polyllysine-coated plates and further incubated for 18-24 h in the presence of differentiating buffer before use.

The Ca^{2+} Kit Assay (Molecular Devices), based on a fluorescent calcium indicator 485-535 nm wavelength, was used.

Differentiated cells were incubated with dye loading for 30 min at 37°C then, nifedipine alone (1 μ M) or in the presence of ω -conotoxin or test compounds were added for further 15 min.

The fluorescence (485-535nm) was measured before and after (30-40 sec) the automated injection of 100 mM KCl depolarizing solution using a Victor plate reader (Perkin Elmer).

Example 11

Mice Formalin Test

According to a modified protocol from Rosland et al., (1990) mice were

injected subcutaneously (s.c.) with 20 μ l of 2.7% solution of formalin into the plantar surface of left hindpaw and placed immediately into clear PVC observation chambers (23 x 12 x 13 cm). Pain behaviour was quantified by counting the cumulative licking time (s) of the injected paw. Measurements
5 were taken during the early phase (0-5 min) and late phase (30-40 min) after formalin injection. (Tjolsen et al. 1992).

The test compound was administered p.o. 15 min before formalin injection in a volume of 10 ml/kg body weight to groups of 10 mice per dose. Control group was treated with vehicle.

10 Example 12


Carragenan model of inflammation

Male Wistar rats of 175-200 grams were used.

The left hind paw was injected with 100 μ l of carrageenan (2% w/v in saline). Compounds of the invention (30 mg/kg), indomethacin (5 mg/kg) or
15 control vehicle (such as distilled water) were orally administered 1 h before carrageenan injection. The paw volume was measured with a plethysmometer (Ugo Basile) immediately before (basal) and 1, 2, 3, 4 and 5h after the carrageenan injection.

CLAIMS

1. The use of a compound of general formula I

**I**

wherein

m is an integer from 1 to 3

X is methylene, oxygen, sulphur or a NR^6 group;

10 **R¹** is a straight or branched $\text{C}_1\text{-C}_8$ alkyl or $\text{C}_3\text{-C}_8$ alkenylene or $\text{C}_3\text{-C}_8$ alkynylene chain, optionally substituted with CF_3 , phenyl, phenoxy or naphthyl, the aromatic rings optionally substituted by one or more $\text{C}_1\text{-C}_4$ alkyl, halogens, trifluoromethyl, hydroxy or $\text{C}_1\text{-C}_4$ alkoxy groups;

15 **R², R³** are independently hydrogen, a $\text{C}_1\text{-C}_3$ alkyl chain, halogen, trifluoromethyl, hydroxy or $\text{C}_1\text{-C}_4$ alkoxy groups;

R⁴, R⁵, R⁶ are independently hydrogen or $\text{C}_1\text{-C}_6$ alkyl;

and the pharmaceutically acceptable salts thereof for the preparation of a medicament having sodium and/or calcium channel modulating activity.

2. A compound of general formula I as specified in claim 1, provided that
20 when **R¹** is phenyl, benzyl, 2-phenethyl or 3-phenpropyl optionally and independently substituted on the phenyl ring by one or two $\text{C}_1\text{-C}_6$ alkyl, halogen, hydroxy, $\text{C}_1\text{-C}_4$ alkoxy or trifluoromethyl and **X** is oxygen, sulphur, methylene or -NH- , at least one of **R²** or **R³** is other than hydrogen.

3. The use of a compound according to claim 1 having the general formula
25 **I** as specified in claim 1, wherein **m** is 1 or 2, **X** is oxygen or methylene or NH

or NCH_3 , R^1 is $\text{C}_1\text{-C}_8$ alkyl chain, optionally substituted with CF_3 , phenyl or phenoxy group, where the aromatic ring in R^1 is optionally substituted by one or two halogen or methoxy or trifluoromethyl groups, R^2 and R^3 are hydrogen, methyl, methoxy, fluorine, chlorine or bromine, R^4 and R^5 are hydrogen or methyl, halogen is chlorine or fluorine.

4. The use according to claim 1 wherein the compound is selected from the group consisting of:

- 3-(4-Butyloxy-benzylamino)-pyrrolidin-2-one;
- 3-[4-(4-trifluorobutyloxy)-benzylamino]-pyrrolidin-2-one;
- 10 3-(4-Pentyloxy-benzylamino)-pyrrolidin-2-one;
- 3-[4-(5-trifluoropentyloxy)-benzylamino]-pyrrolidin-2-one;
- 3-(4-Phenylethyl-benzylamino)-pyrrolidin-2-one;
- 3-(4-Benzyloxy-benzylamino)-pyrrolidin-2-one;
- 3-(4-Phenylbutoxy-benzylamino)-pyrrolidin-2-one;
- 15 3-(4-Phenylpentyloxy-benzylamino)-pyrrolidin-2-one;
- 3-(4-Phenylallyloxy)-benzylamino-pyrrolidin-2-one;
- 3-(4-Phenoxyethoxy-benzylamino)-pyrrolidin-2-one;
- 3-[4-(Naphthalen-1-ylmethoxy)-benzylamino]-pyrrolidin-2-one;
- 3-(4-Pentyloxy-3-fluoro-benzylamino)-pyrrolidin-2-one;
- 20 3-(4-Pentyloxy-3-chloro-benzylamino)-pyrrolidin-2-one;
- 3-(4-Pentyloxy-3-bromo-benzylamino)-pyrrolidin-2-one;
- 3-(4-Pentyloxy-3-methoxy-benzylamino)-pyrrolidin-2-one;
- 3-(4-Pentyloxy-3-methyl-benzylamino)-pyrrolidin-2-one;
- 3-(4-Benzyloxy-3-fluoro-benzylamino)-pyrrolidin-2-one;
- 25 3-(4-Benzyloxy-3-bromo-benzylamino)-pyrrolidin-2-one;
- 3-(4-Benzyloxy-3-methoxy-benzylamino)-pyrrolidin-2-one;
- 3-(4-Benzyloxy-3-methyl-benzylamino)-pyrrolidin-2-one;
- 3-(4-Phenylpentyloxy-2-chloro-benzylamino)-pyrrolidin-2-one;

- 3-(4-Phenylpentoxy-3-bromo-benzylamino)-pyrrolidin-2-one;
3-(4-Phenylpentoxy-3-methoxy-benzylamino)-pyrrolidin-2-one;
3-(4-Phenylpentoxy-3-methyl-benzylamino)-pyrrolidin-2-one;
3-(4-Phenylallyloxy-2-chloro-benzylamino)-pyrrolidin-2-one;
5 3-(4-Phenylallyloxy-3-fluoro-benzylamino)-pyrrolidin-2-one;
3-(4-Phenylallyloxy-3-bromo-benzylamino)-pyrrolidin-2-one;
3-(4-Phenylallyloxy-3-methoxy-benzylamino)-pyrrolidin-2-one;
3-(4-Phenylallyloxy-3-methyl-benzylamino)-pyrrolidin-2-one;
3-(4-Phenoxyethoxy-2-chloro-benzylamino)-pyrrolidin-2-one;
10 3-(4-Phenoxyethoxy-3-fluoro-benzylamino)-pyrrolidin-2-one;
3-(4-Phenoxyethoxy-3-bromo-benzylamino)-pyrrolidin-2-one;
3-(4-Phenoxyethoxy-3-methoxy-benzylamino)-pyrrolidin-2-one;
3-(4-Phenoxyethoxy-3-methyl-benzylamino)-pyrrolidin-2-one;
3-[4-(Naphthalen-1-ylmethoxy)-3-bromo-benzylamino]-pyrrolidin-2-one;
15 3-[4-(Naphthalen-1-ylmethoxy)-3-methoxy-benzylamino]-pyrrolidin-2-one;
3-[4-(Naphthalen-1-ylmethoxy)-3-methyl-benzylamino]-pyrrolidin-2-one;
3-(4-Pentyloxy-3-bromo-5-methoxy-benzylamino)-pyrrolidin-2-one;
3-(4-Pentyloxy-3,5-dimethoxy-benzylamino)-pyrrolidin-2-one;
3-(4-Pentyloxy-3,5-dimethyl-benzylamino)-pyrrolidin-2-one;
20 3-(4-Benzyloxy-3-bromo-5-methoxy-benzylamino)-pyrrolidin-2-one;
3-(4-Benzyloxy-3,5-dimethoxy-benzylamino)-pyrrolidin-2-one;
3-(4-Benzyloxy-3,5-dimethyl-benzylamino)-pyrrolidin-2-one;
3-(4-Phenylallyloxy-3-bromo-5-methoxy-benzylamino)-pyrrolidin-2-one;
3-(4-Phenylallyloxy-3,5-dimethoxy-benzylamino)-pyrrolidin-2-one;
25 3-(4-Phenylallyloxy-3,5-dimethyl-benzylamino)-pyrrolidin-2-one;
3-(4-Phenylpentoxy-3-bromo-5-methoxy-benzylamino)-pyrrolidin-2-one;
3-(4-Phenylpentoxy-3,5-dimethoxy-benzylamino)-pyrrolidin-2-one;
3-(4-Phenylpentoxy-3,5-dimethyl-benzylamino)-pyrrolidin-2-one;

- 3-(4-Phenoxyethoxy-3-bromo-5-methoxy-benzylamino)-pyrrolidin-2-one;
3-(4-Phenoxyethoxy-3,5-dimethoxy-benzylamino)-pyrrolidin-2-one;
3-(4-Phenoxyethoxy-3,5-dimethyl-benzylamino)-pyrrolidin-2-one;
3-[4-(Naphthalen-1-ylmethoxy)-2-chloro-5-methoxy-benzylamino]-
5 pyrrolidin-2-one;
3-[4-(Naphthalen-1-ylmethoxy)-3-fluoro-5-methoxy-benzylamino]-
pyrrolidin-2-one;
3-[4-(Naphthalen-1-ylmethoxy)-3-bromo-5-methoxy-benzylamino]-
pyrrolidin-2-one;
10 3-[4-(Naphthalen-1-ylmethoxy)-3,5-dimethoxy-benzylamino]-
pyrrolidin-2-one;
3-[4-(Naphthalen-1-ylmethoxy)-3,5-dimethyl-benzylamino]-pyrrolidin-
2-one;
3-[4-(2-Fluorobenzyloxy)-benzylamino]-pyrrolidin-2-one;
15 3-[4-(2-Fluorobenzyloxy)-benzylamino]-N-methylpyrrolidin-2-one;
3-[4-(2-triFluoromethyl-benzyloxy)-benzylamino]-pyrrolidin-2-one;
3-[4-(2-Chlorobenzyloxy)-benzylamino]-pyrrolidin-2-one;
3-[4-(2-Methoxybenzyloxy)-benzylamino]-pyrrolidin-2-one;
3-[4-(3-Fluorobenzyloxy)-benzylamino]-pyrrolidin-2-one;
20 3-[4-(3-Fluorobenzyloxy)-benzylamino]-N-methylpyrrolidin-2-one;
3-{N-[4-(3-Fluorobenzyloxy)-benzyl]-N-methyl}-amino-pyrrolidin-2-one;
3-[4-(3-triFluoromethyl-benzyloxy)-benzylamino]-pyrrolidin-2-one;
3-[4-(3-Chlorobenzyloxy)-benzylamino]-pyrrolidin-2-one;
3-[4-(3-Methoxybenzyloxy)-benzylamino]-pyrrolidin-2-one;
25 3-[4-(3-Methoxybenzyloxy)-benzylamino]-N-methylpyrrolidin-2-one;
3-[4-(4-Fluorobenzyloxy)-benzylamino]-pyrrolidin-2-one;
3-[4-(4-Chlorobenzyloxy)-benzylamino]-pyrrolidin-2-one;
3-[4-(4-Methoxybenzyloxy)-benzylamino]-pyrrolidin-2-one;

- 3-[4-(4-triFluoromethyl-benzyloxy)-benzylamino]-pyrrolidin-2-one;
3-[4-(2,3-diChlorobenzyloxy)-benzylamino]-pyrrolidin-2-one;
3-[4-(3,4-diChlorobenzyloxy)-benzylamino]-pyrrolidin-2-one;
3-[4-(3,4-diMethoxybenzyloxy)-benzylamino]-pyrrolidin-2-one;
5 3-[4-(3,5-diMethoxybenzyloxy)-benzylamino]-pyrrolidin-2-one;
3-[4-(3,5-diMethoxybenzyloxy)-benzylamino]-N-methylpyrrolidin-2-one;
3-[4-(3,5-diMethoxyphenyl)-pentoxy]-benzylamino-pyrrolidin-2-one;
3-[4-(2-Fluorobenzyloxy)-3-methyl-benzylamino]-pyrrolidin-2-one;
3-[4-(2-triFluoromethyl-benzyloxy)-3-methyl-benzylamino]-pyrrolidin-
10 2-one;
3-[4-(3-Fluorobenzyloxy)-3-methyl-benzylamino]-pyrrolidin-2-one;
3-{[4-(3-Fluorobenzyloxy)-3-methyl-benzyl]-N-methylamino}-
pyrrolidin-2-one;
3-[4-(3-triFluoromethyl-benzyloxy)-3-methyl-benzylamino]-pyrrolidin-
15 2-one;
3-[4-(3-Chlorobenzyloxy)-3-methyl-benzylamino]-pyrrolidin-2-one;
3-{[4-(3-Chlorobenzyloxy)-3-methyl-benzyl]-N-methylamino}-
pyrrolidin-2-one;
3-[4-(3-Bromobenzyloxy)-3-methyl-benzylamino]-pyrrolidin-2-one;
20 3-{[4-(3-Bromobenzyloxy)-3-methyl-benzyl]-N-methylamino}-
pyrrolidin-2-one;
3-[4-(4-triFluoromethyl-benzyloxy)-2-chloro-benzylamino]-pyrrolidin-
2-one;
3-[4-(4-Fluorobenzyloxy)-3-methyl-benzylamino]-pyrrolidin-2-one;
25 3-[4-(4-triFluoromethyl-benzyloxy)-3-fluoro-benzylamino]-pyrrolidin-
2-one;
3-[4-(4-triFluoromethyl-benzyloxy)-3-bromo-benzylamino]-pyrrolidin-
2-one;

- 3-[4-(4-triFluoromethyl-benzyloxy)-3-methoxy-benzylamino]-pyrrolidin-2-one;
- 3-[4-(4-triFluoromethyl-benzyloxy)-3-methyl-benzylamino]-pyrrolidin-2-one;
- 5 3-[4-(4-Chlorobenzyloxy)-3-methyl-benzylamino]-pyrrolidin-2-one;
- 3-[4-(4-triFluoromethyl-benzyloxy)-3-bromo-5-methoxy-benzylamino]-pyrrolidin-2-one;
- 3-[4-(4-triFluoromethyl-benzyloxy)-3,5-dimethoxy-benzylamino]-pyrrolidin-2-one;
- 10 3-[4-(4-triFluoromethyl-benzyloxy)-3,5-dimethyl-benzylamino]-pyrrolidin-2-one;
- 3-[4-(3,4-diChlorobenzyloxy)-2-chloro-benzylamino]-pyrrolidin-2-one;
- 3-[4-(3,4-diChlorobenzyloxy)-3-fluoro-benzylamino]-pyrrolidin-2-one;
- 3-[4-(3,4-diChlorobenzyloxy)-3-bromo-benzylamino]-pyrrolidin-2-one;
- 15 3-[4-(3,4-diChlorobenzyloxy)-3-methoxy-benzylamino]-pyrrolidin-2-one;
- 3-[4-(3,4-diChlorobenzyloxy)-3-methyl-benzylamino]-pyrrolidin-2-one;
- 3-[4-(3,5-diMethoxybenzyloxy)-2-chloro-benzylamino]-pyrrolidin-2-one;
- 3-[4-(3,5-diMethoxybenzyloxy)-3-fluoro-benzylamino]-pyrrolidin-2-one;
- 3-[4-(3,5-diMethoxybenzyloxy)-3-bromo-benzylamino]-pyrrolidin-2-one;
- 20 3-[4-(3,5-diMethoxybenzyloxy)-3-methoxy-benzylamino]-pyrrolidin-2-one;
- 3-[4-(3,5-diMethoxybenzyloxy)-3-methyl-benzylamino]-pyrrolidin-2-one;
- 3-[4-(3,4-diChlorobenzyloxy)-3,5-dimethoxy-benzylamino]-pyrrolidin-2-one;
- 3-[4-(3,4-diChlorobenzyloxy)-3,5-dimethyl-benzylamino]-pyrrolidin-2-one;
- 25 3-[4-(3,5-diChlorobenzyloxy)-3-bromo-5-methoxy-benzylamino]-pyrrolidin-2-one;
- 3-[4-(3,5-diMethoxybenzyloxy)-3-bromo-5-methoxy-benzylamino]-pyrrolidin-2-one;

36

3-[4-(3,5-diMethoxybenzyloxy)-3,5-dimethoxy-benzylamino]-pyrrolidin-2-one;

3-[4-(3,5-diMethoxyphenyl)-allyloxy)-3,5-dimethoxy-benzylamino]-pyrrolidin-2-one.

5 3-(4-Benzyloxy-benzylamino)-piperidin-2-one;

3-(4-Benzyloxy-benzylamino)-azepan-2-one;

3-[4-(2-Fluorobenzyloxy)-benzylamino]-piperidin-2-one;

3-[4-(2-Fluorobenzyloxy)-benzylamino]-azepan-2-one;

3-[4-(2-Chlorobenzyloxy)-benzylamino]-piperidin-2-one;

10 3-[4-(2-Chlorobenzyloxy)-benzylamino]-azepan-2-one;

3-[4-(3-Fluorobenzyloxy)-benzylamino]-piperidin-2-one;

3-[4-(3-Fluorobenzyloxy)-benzylamino]-azepan-2-one;

3-[4-(4-Fluorobenzyloxy)-benzylamino]-piperidin-2-one;

3-[4-(4-Fluorobenzyloxy)-benzylamino]-azepan-2-one;

15 3-[4-(2-Chlorobenzylamino)-benzylamino]-piperidin-2-one;


3-[4-(2-Chlorobenzylamino)-benzylamino]-azepan-2-one;

3-{4-[(2-Chlorobenzyl)methylamino]-benzylamino}-piperidin-2-one;

3-{4-[(2-Chlorobenzyl)methylamino]-benzylamino}-azepan-2-one;

or pharmaceutically acceptable salts thereof.

- 20 5. A process for the preparation of a compound of formula I, as defined in claim 1, or a pharmaceutically acceptable salt thereof, the process comprising:
a) reaction of compounds of formula II




II

25

37

wherein R^1 , R^2 , R^3 and X are as defined above


with compounds of formula III, in the presence of a reducing agent



III

wherein m and R^5 are as defined previously thus obtaining a compound of formula I; or

b) reaction of compounds of formula IV




IV

wherein R^1 , R^2 , R^3 and X are as defined above and Y is a halogen atom or a O-EWG group, where the EWG means an electron withdrawing group, like e.g. mesyl, tosyl or trifluoroacetyl groups, able to transform the oxygen which

they are linked to, in a good leaving group with compounds of formula III thus obtaining a compound of formula I; or

c) reacting of a compound of formula V



V

38

wherein R^1 , R^2 , R^3 , R^5 , X and m are as defined above, with compounds of formula VI or VII



VI




VII

wherein Y is as defined above; R^4 is as above defined and R^7 is hydrogen or C_1-C_5 alkyl; and, if desired, converting a compound of the invention into another compound of the invention and/or, if desired, converting a compound of the invention into a pharmaceutically acceptable salt and/or, if desired, converting a salt into a free compound and/or, if desired, separating a mixture of isomers of compounds of the invention into a single isomer.

6. A pharmaceutical composition containing a compound of formula I, as defined in claim 1, or a pharmaceutically acceptable salt thereof, in admixture with a suitable carrier and/or diluent and optionally to other therapeutic agents.

ABSTRACT**3-AMINOPYRROLIDONE DERIVATIVES**

5 This invention is related to compounds and use of 3-aminopyrrolidone derivatives and analogues of the following general formula (I)



I

10 wherein

m is an integer from 1 to 3

X is methylene, oxygen, sulphur or a NR^6 group;

R¹ is a straight or branched $\text{C}_1\text{-C}_8$ alkyl or $\text{C}_3\text{-C}_8$ alkenylene or $\text{C}_3\text{-C}_8$ alkynylene chain, optionally substituted with CF_3 , phenyl, phenoxy or naphthyl, the aromatic rings optionally substituted by one or more $\text{C}_1\text{-C}_4$ alkyl, halogens, trifluoromethyl, hydroxy or $\text{C}_1\text{-C}_4$ alkoxy groups;

R², R³ are independently hydrogen, a $\text{C}_1\text{-C}_3$ alkyl chain, halogen, trifluoromethyl, hydroxy or $\text{C}_1\text{-C}_4$ alkoxy groups;

20 **R⁴, R⁵, R⁶** are independently hydrogen or $\text{C}_1\text{-C}_6$ alkyl;

and the pharmaceutically acceptable salts thereof that are active as sodium and/or calcium channel modulators and therefore useful in preventing, alleviating and curing a wide range of pathologies, including, but not limited to, neurological, psychiatric, cardiovascular, inflammatory, ophthalmic, urologic, metabolic and gastrointestinal diseases, where the above mechanisms have been described as playing a pathological role.

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/EP04/012957

International filing date: 16 November 2004 (16.11.2004)

Document type: Certified copy of priority document

Document details: Country/Office: EP
Number: 03026779.3
Filing date: 21 November 2003 (21.11.2003)

Date of receipt at the International Bureau: 03 February 2005 (03.02.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse

This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record.

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☒ **BLACK BORDERS**

☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☐ **FADED TEXT OR DRAWING**

☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☐ **SKEWED/SLANTED IMAGES**

☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☐ **GRAY SCALE DOCUMENTS**

☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☒ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.